

Homeomorphisms of $\overline{U} \times R$ and rotation number

Paul Fabel
 Drawer MA
 Mississippi State University
 fabel@ra.msstate.edu

Abstract

Suppose $U \subset R^2$ is bounded open and contractible and $H : \overline{U} \times R \rightarrow \overline{U} \times R$ is a homeomorphism leaving invariant $U \times R$. If ∂U is locally connected and not a simple closed curve H induces a homeomorphism of the solid cylinder leaving invariant sufficiently many vertical lines to determine a rotation number. If ∂U is not locally connected H admits a natural notion of rotation number despite a general absence of an induced homeomorphism of the solid cylinder.

1 Introduction

The following theorem, credited to Ursell and Young [11] is central to applications ([1],[3],[2],[4],[5],[9]) of prime end theory to dynamical systems:

Theorem 1 *Suppose $U \subset R^2$ is bounded open and contractible, $\psi : U \rightarrow \text{int}(D^2)$ is conformal, and $h : \overline{U} \rightarrow \overline{U}$ is a homeomorphism such that $h(U) = U$. Then $\psi h \psi^{-1}$ can be extended to a homeomorphism of D^2 .*

In particular if h is orientation preserving then $\overline{\psi h \psi^{-1}}_{S^1} : \partial D^2 \rightarrow \partial D^2$ determines a rotation number, measuring in a sense the average rotation by h of ∂U about U .

Specific examples of higher dimensional versions of Theorem 1 are in short supply despite a general criteria established in [4].

For example if ∂U is not locally connected, there is no corresponding version of Theorem 1 for domains $U \times R \subset R^3$ with $\Psi : U \times R \rightarrow \text{int}(D^2) \times R$ defined via $\psi(u, t) = (\psi(u), t)$. Example 4 exhibits a homeomorphism $H : \overline{U} \times R \rightarrow \overline{U} \times R$ such that $H(U \times R) = U \times R$ but $\Psi H \Psi^{-1}$ **cannot** be extended to a homeomorphism of $D^2 \times R$.

The main result of this paper (Theorem 18) salvages a notion of rotation number for such homeomorphisms $H : \overline{U} \times R \rightarrow \overline{U} \times R$. The basic idea is that H preserves the circular order of a certain collection of sets, each of which can be understood as the product of R with an interval of accessible prime ends of U . If ∂U is not locally connected this provides enough information to determine a homeomorphism $g : S^1 \rightarrow S^1$ whose rotation number we declare to be that of H .

On the other hand a useful ‘3 dimensional prime end theory’ exists if ∂U is locally connected. Theorem 13 shows Ψ induces a homeomorphism of the (3 cell) two point compactification of $D^2 \times R$. Moreover if ∂U is not a simple closed curve of the cutpoints of ∂U help to determine a discrete collection of invariant boundary lines which in turn determine a rotation number.

Both notions of rotation number are invariant under topological conjugacy, and agree with the usual rotation number of h in the special case $H(u, t) = (h(u), t)$ where $h : \overline{U} \rightarrow \overline{U}$ is a homeomorphism such that $h(U) = U$.

2 Preliminaries

Suppose throughout this paper $U \subset R^2$ is bounded open and contractible, ∂U denotes $\overline{U} \setminus U$, $D^2 \subset R^2$ denotes the closed unit disk and $\psi : U \rightarrow \text{int}(D^2)$ is conformal. All function spaces will have the compact open topology.

Define $\Psi : U \times R \rightarrow \text{int}(D^2) \times R$ via $\Psi(u, t) = (\psi(u), t)$.

Let $D^3 = \{(x, y, z) \in R^3 \mid \sqrt{x^2 + y^2 + z^2} \leq 1\}$, the standard 3 cell.

If $J \subset S^1$ is connected let $\text{int}(J) = J$ if $J = \{x\}$. Otherwise let $\text{int}(J)$ denote the union of all open intervals contained in J .

If Y is a topological space attach two points $\{\infty\}$ and $\{-\infty\}$ to $Y \times R$ creating $(Y \times J) \cup \{\infty, -\infty\}$ topologized such that $(y_n, t_n) \rightarrow \{-\infty\}$ iff $t_n \rightarrow \infty$ and $(y_n, t_n) \rightarrow \{\infty\}$ iff $t_n \rightarrow -\infty$.

In similar fashion we attach two points $\{\infty\}$ and $\{-\infty\}$ to $Y \times (-1, 1)$ and create a new space $\overline{Y \times (-1, 1)}$ topologized such that $(y_n, t_n) \rightarrow \{-\infty\}$ iff $t_n \rightarrow -1$ and $(y_n, t_n) \rightarrow \{\infty\}$ iff $t_n \rightarrow 1$.

3 The map $\phi : X^* \rightarrow D^2$

The following procedure creates a complete metric space X^* whose underlying set can be seen as the union of U and the accessible prime ends of U .

Define a metric $d^* : U \times U \rightarrow R$ such that $d^*(x, y) < \varepsilon$ iff there exists a map $f : [0, 1] \rightarrow U$ such that $f(0) = x, f(1) = y$ and $\forall s, t \mid f(t) - f(s) \mid < \varepsilon$. Let (X^*, d^*) denote the metric completion of (U, d^*) .

Let $\partial X^* = X^* \setminus U^*$.

Lemma 2 *There exists a map $\phi : X^* \rightarrow D^2$ such that ϕ is uniformly continuous, one to one and $\phi(U^*) = \text{int}(D^2)$.*

Proof. Let $\overline{id} : X^* \rightarrow \overline{U}$ denote the unique extension of the uniformly continuous identity map $id : U^* \rightarrow U$. Uniform continuity of $\psi(id)$ is essentially a consequence of Theorem 1 and is proved in Theorem 3.1 of [7]. Define $\phi : X^* \rightarrow D^2$ to be the unique continuous extension of $\psi(id)$. Suppose $x \neq y$ and $\{x, y\} \subset X^*$. If $\overline{id}(x) \neq \overline{id}(y)$ it follows from Proposition 2.14 [10] that $\psi(\overline{id}(x)) \neq \psi(\overline{id}(y))$. If $\overline{id}(x) = \overline{id}(y)$ then $\{\overline{id}(x), \overline{id}(y)\} \subset \partial U$ and $\{x, y\} \subset \partial X^*$. Construct a closed topological disk $E \subset \overline{U}$ such that $\overline{id}(x) \in \partial E$ and $E \setminus \{\overline{id}(x)\} \subset U$. If $d^*(x, y) \neq 0$ then $\text{int}(E) \cap \partial U \neq \emptyset$ and $\overline{id}(x)$ and $\overline{id}(y)$ determine distinct prime ends. Let $z \in \partial E \setminus \{\overline{id}(x)\}$. By Theorem 2.15 [10] ψ determines a bijection between the prime ends of U and ∂D^2 . In particular $\partial E \setminus \{z\}$ determines distinct endcuts which map under ψ to distinct points of ∂D^2 . Hence ϕ is one to one. ■

Remark 3 *The injective map $\phi : X^* \rightarrow D^2$ need not be an embedding. For example if $U \subset R^2$ is the region bounded by a ‘Warsaw circle’. The canonical map from $\partial X^* \rightarrow \partial D^2$ is a continuous bijection but not a homeomorphism.*

4 Failure of homeomorphism extension

Example 4 *Suppose $U \subset R^2$ is the interior of the standard Warsaw circle. There there exists $H \in G(U \times R)$ such that $\Psi H \Psi^{-1} : \text{int}(D^2) \times R \rightarrow \text{int}(D^2) \times R$ is not extendable to a homeomorphism of $D^2 \times R$.*

Proof. Let $U \subset R^2$ be the interior of a standardly embedded Warsaw circle such that the impression of the bad prime end (the closed interval ‘limit bar’) is precisely $\partial U \cap (\{0\} \times R)$. Define a homeomorphism $H : U \times R \rightarrow U \times R$ via $H(x, y, t) = (x, y, y + t)$. Note $\Psi H \Psi^{-1}$ is not continuously extendable near the bad prime end since H is not level preserving on the impression of the bad prime end. ■

5 Homeomorphisms of S^1 and rotation number

The notion of rotation number of a homeomorphism of the unit circle dates back to Poincare. Its properties are derived in Devaney’s book [6] formally for diffeomorphisms but the proofs are valid for homeomorphisms. See also [8] for a helpful survey.

Theorem 5 Suppose $g : S^1 \rightarrow S^1$ is an orientation preserving homeomorphism, $\Pi : R \rightarrow S^1$ is the covering map $\Pi(\theta) = e^{2\pi i\theta}$, $x \in R$ and $G_1, G_2 : R \rightarrow R$ are homeomorphisms such that $\Pi(G_i) = g\Pi$. Then the following limits exist and differ by an integer: $\lim_{n \rightarrow \infty} \frac{G_1^n(x)}{n}$ and $\lim_{n \rightarrow \infty} \frac{G_2^n(x)}{n}$. This number (mod 1) is the **rotation number** of g and is invariant under the choice of x . If g reverses orientation, then g has two fixed points and we declare g to have rotation number 0. The rotation number of g is invariant under topological conjugacy (if $h : S^1 \rightarrow S^1$ is a homeomorphism and $\hat{g} = hgh^{-1}$ then $\text{rot}(\hat{g}) = \text{rot}(g)$). Finally, rot is a continuous function on the space of homeomorphisms of S^1 .

6 Sets with circular order and rotation number

Given 4 distinct point $\{x_1, x_2, x_3, x_4\} \subset S^1$ declare $x_1 < x_2 < x_3 < x_4$ if there exists a homeomorphism $g : S^1 \rightarrow S^1$ such that $g(i^n) = x_n$.

Note it is allowed that g reverses orientation. Suppose $\{J_1, J_2, J_3, J_4\}$ is a collection of distinct pairwise disjoint nonempty subsets of S^1 . Declare $J_1 < J_2 < J_3 < J_4$ if $x_1 < x_2 < x_3 < x_4$ whenever $x_i \in J_i$.

Suppose J is a proper connected subset of S^1 . Define $\text{int}(J) = J$ if $J = \{x\}$ and define $\text{int}(J)$ to be the largest open interval contained in J otherwise.

Definition 6 Suppose $\hat{A}$ is a collection pairwise disjoint subsets of S^1 and $h : \hat{A} \rightarrow \hat{A}$ is a bijection. Then h is order preserving if there exists a homeomorphism $g : S^1 \rightarrow S^1$ such that $g(J) = h(J)$ for each element $J \in \hat{A}$. The homeomorphism g is said to be **compatible** with h . If $|\hat{A}| \geq 3$ then declare h orientation preserving/reversing iff g is orientation preserving/reversing.

Lemma 7 Suppose $\hat{A}$ is a collection pairwise disjoint subsets of S^1 such that $|\hat{A}| \geq 4$ and $\cup_{J \in \hat{A}} J$ is dense in S^1 . Suppose $J = \text{int}(J)$ for each $J \in \hat{A}$. Suppose $h : \hat{A} \rightarrow \hat{A}$ is a bijection leaving invariant the set of nontrivial elements of $\hat{A}$. Then h is order preserving if and only if $h(J_3)$ and $h(J_4)$ belong to the same component of $S^1 \setminus \{h(J_1) \cup h(J_2)\}$ whenever $J_1 < J_2 < J_3 < J_4$.

Proof. Suppose h is order preserving. Let $g : S^1 \rightarrow S^1$ be a homeomorphism such that $g(J) = h(J)$ for each $J \in \hat{A}$. Suppose $J_1 < J_2 < J_3 < J_4$. Then J_3 and J_4 belong to the same component of $S^1 \setminus \{J_1 \cup J_2\}$. Since g is a homeomorphism $g(J_3)$ and $g(J_4)$ belong to the same component of $S^1 \setminus \{g(J_1) \cup g(J_2)\}$. The conclusion follows since $g(J_i) = h(J_i)$. Conversely suppose $h(J_3)$ and $h(J_4)$ belong to the

same component of $S^1 \setminus \{h(J_1) \cup h(J_2)\}$ whenever $J_1 < J_2 < J_3 < J_4$. Suppose $J_1 < J_2 < J_3 < J_4$. Then either $h(J_1) < h(J_2) < h(J_3) < h(J_4)$ or $h(J_1) < h(J_2) < h(J_4) < h(J_3)$. Suppose in order to obtain a contradiction that $h(J_1) < h(J_2) < h(J_4) < h(J_3)$. Then $h(J_2)$ and $h(J_3)$ lie in opposite components of $S^1 \setminus \{h(J_1) \cup h(J_4)\}$. On the other hand by hypothesis $J_4 < J_1 < J_2 < J_3$ and hence $h(J_2)$ and $h(J_3)$ lie in the same component of $S^1 \setminus \{h(J_1) \cup h(J_4)\}$ and we have a contradiction. Thus $h(J_1) < h(J_2) < h(J_3) < h(J_4)$ and globally the bijection h either preserves or reverses orientation. Let $g : S^1 \rightarrow S^1$ be the unique homeomorphism mapping J linearly onto $h(J)$ whenever J is a nontrivial component of $\hat{A}$. ■

Lemma 8 *Suppose $\hat{A}$ is a collection of pairwise disjoint connected subsets of S^1 such that $|\hat{A}| \geq 3$ and $\cup_{J \in \hat{A}} J$ is dense in S^1 . Suppose $J = \text{int}(J)$ whenever $J \in \hat{A}$ and suppose $h : \hat{A} \rightarrow \hat{A}$ is an order preserving bijection as demonstrated by the compatible orientation preserving homeomorphisms $g_1 : S^1 \rightarrow S^1$ and $g_2 : S^1 \rightarrow S^1$. Then g_1 and g_2 have the same rotation number.*

Proof. Let $B \subset S^1$ denote the complement of the union of the open intervals of $\hat{A}$. Then $B \neq \emptyset$, $g_1(B) = B$ and $g_2|_B = g_1|_B$. Thus there exists $\tilde{b} \in B$ and lifts $G_1 : R \rightarrow R$ and $G_2 : R \rightarrow R$ respectively of g_1 and g_2 such that $\pi(\tilde{b}) \in B$ and $G_1^n(\tilde{b}) = G_2^n(\tilde{b}) \forall n$. Hence by Theorem 5 g_1 and g_2 have the same rotation number. ■

7 The order preserving bijection $h : \hat{A} \rightarrow \hat{A}$

Define a surjection $p : D^2 \times [-1, 1] \rightarrow D^3$ via $p(x, y, t) = ((1 - |t|)x, (1 - |t|)y, t)$. Note p collapses $D^2 \times \{-1\}$ and $D^2 \times \{1\}$ to points and is otherwise injective.

Note the map $\Phi_{X^* \times (-1, 1)} : X^* \times (-1, 1) \rightarrow D^2 \times (-1, 1)$ defined such that $\Phi(x, t) = (\phi(x), t)$ can be continuously extended to an injective map $\Phi : \overline{X^* \times (-1, 1)} \hookrightarrow \overline{D^2 \times (-1, 1)}$ such that $\Phi(\infty) = \infty$ and $\Phi(-\infty) = -\infty$.

Define a metric on $X^* \times R$ via $d((x, t), (y, t)) = \max(d^*(x, y), |s - t|)$.

Define a set $\hat{A} \subset 2^{\partial D^2}$ such that $\beta \in \hat{A}$ iff $\beta = \text{int}(\phi(X))$ for some path component $X \subset \partial X^*$.

Theorem 9 *The homeomorphism $H : \overline{U} \times R \rightarrow \overline{U} \times R$ induces a canonical homeomorphism $H^* : X^* \times R \rightarrow X^* \times R$ such that $H^*(\partial X^* \times R) = \partial X^* \times R$ and such that H^* is extendable to a homeomorphism of $(X^* \times R) \cup \{\infty, -\infty\}$.*

Proof. We first obtain, as follows, an induced homeomorphism $H^* : X^* \times R \rightarrow X^* \times R$ such that $H^*(U^* \times R) = U^* \times R$ and for each S and T there exists $S^\wedge$ and $T^\wedge$ such $H^*(X^* \times [S, T]) \cup H^{*-1}(X^* \times [S, T]) \subset X^* \times [S^\wedge, T^\wedge]$. Suppose $\{(z_n, t_n)\}$ is Cauchy in $U^* \times R$. Since $\{t_n\}$ is Cauchy choose S and T such that $\forall n (z_n, t_n) \in X^* \times [S, T]$. Moreover for each n, m we may choose a path $\gamma_{nm} \subset U^* \times [S, T]$ such that γ_{nm} connects (z_n, t_n) to (z_m, t_m) and $\text{diam}(\gamma_{nm}) < 2d((z_n, t_n), (z_m, t_m))$. Hence $\lim_{n,m \rightarrow \infty} \text{diam}(\gamma_{nm}) = 0$. However, since $\overline{U} \times [S, T]$ is compact, both $H_{\overline{U} \times [S, T]}$ and $H_{\overline{U} \times [S, T]}^{-1}$ are uniformly continuous. Thus $\lim_{n,m \rightarrow \infty} \text{diam}(H(\gamma_{nm})) = 0 = \lim_{n,m \rightarrow \infty} \text{diam}(H^{-1}(\gamma_{nm}))$. Hence H and H^{-1} preserve Cauchy sequences in $U^* \times R$ and thus are extendable to maps $H^* : X^* \times R \rightarrow X^* \times R$ and $(H^{-1})^* : X^* \times R \rightarrow X^* \times R$ such that $H^*((H^{-1})^*) = (H^{-1})^*H = id$. Since id is a bijection it follows that H^* and $(H^{-1})^*$ are bijections and hence homeomorphisms. By definition $H(U \times R) = H^{-1}(U \times R) = U \times R$. Since compact subsets of $\overline{U} \times R$ are bounded we may choose $S^\wedge$ and $T^\wedge$ such that the compactum $H(\overline{U} \times [S, T]) \cup H^{-1}(\overline{U} \times [S, T]) \subset \overline{U} \times [S^\wedge, T^\wedge]$. Note $\overline{U} \times [S, T]$ (and hence $H(\overline{U} \times [S, T])$) separate $\overline{U} \times R$ into two components. Since $\overline{U} \times (T, \infty)$ is connected $H(\overline{U} \times (T, \infty))$ intersects and contains exactly one component of $(\overline{U} \times R) \setminus (\overline{U} \times [S^\wedge, T^\wedge])$. Thus if $(z_n, t_n) \rightarrow \infty$ then $H(z_n, t_n)$ converges either to ∞ to $-\infty$. Thus both H^* and $(H^*)^{-1}$ are extendable to maps of $(X^* \times R) \cup \{\infty, -\infty\}$ and both must be homeomorphisms since $H^*(H^{*-1})$ is fixes pointwise a dense set. ■

Corollary 10 *The induced homeomorphism $H^* : X^* \times R \rightarrow X^* \times R$ induces a canonical homeomorphism $H^{**} : \overline{X^* \times (-1, 1)} \rightarrow \overline{X^* \times (-1, 1)}$ and a canonical bijection $h : A^\wedge \rightarrow A^\wedge$.*

Proof. Let $H^* : X^* \times R \rightarrow X^* \times R$ be induced from $H : \overline{U} \times R \rightarrow \overline{U} \times R$ as in Theorem 9. Let $T : R \rightarrow (-1, 1)$ be any homeomorphism. Define a homeomorphism $H^{**} : \overline{X^* \times (-1, 1)} \rightarrow \overline{X^* \times (-1, 1)}$ such that $H^{**}(x, T(t)) = (y, T(s))$ if $H^*(x, t) = (y, s)$. Since $H^{**}(\partial X^* \times (-1, 1)) = \partial X^* \times (-1, 1)$ the homeomorphism H^{**} permutes the path components of $\partial X^* \times (-1, 1)$. Each path component of $\partial X^* \times (-1, 1)$ is of the form $X \times (-1, 1)$ where X is a path component of ∂X^* . By definition $\beta \in A^\wedge$ iff there exists a path component $X \subset \partial X^*$ such that $\beta = \text{int}\phi(X)$. Thus we define a bijection $h : A^\wedge \rightarrow A^\wedge$ satisfying $h(\beta) = \gamma$ if $\beta = \text{int}\phi(X)$ and $H^{**}(X \times (-1, 1)) = Y \times (-1, 1)$ and $\gamma = \text{int}\phi(Y)$. ■

Lemma 11 *Suppose $D \subset D^3$ is a topological disk such that $\text{int}(D) \subset \text{int}(D^3)$, and $\partial D \subset \partial D^3$. Suppose $\{(0, 0, 1), (0, 0, -1)\} \subset \partial D$ and suppose $\alpha : [0, 1] \rightarrow D^3 \setminus D$ satisfies $\alpha(t) \in \text{int}(D^3)$ iff $0 < t < 1$. Sup-*

pose β_1 and β_2 are disjoint nonempty connected subsets of S^1 . Suppose $\partial D \subset p((\beta_1 \cup \beta_2) \times [-1, 1])$. Suppose $\{\alpha(0), \alpha(1)\} \cap p((\beta_1 \cup \beta_2) \times [-1, 1]) = \emptyset$. Then $\pi_1(p^{-1}(\alpha(0), \alpha(1)))$ is contained in a single component of $S^1 \setminus \{\beta_1 \cup \beta_2\}$.

Proof. The topological disk D separates D^3 into two components and ∂D separates ∂D^3 into two components. Moreover the two components of $\partial D^3 \setminus \partial D$ are contained in distinct components of $D^3 \setminus D$. Hence, since $im(\alpha)$ is connected and $im(\alpha) \cap D = \emptyset$, $\partial\alpha$ belongs to a single component of $\partial D^3 \setminus \partial D$. Moreover $\partial D^3 \setminus p((\beta_1 \cup \beta_2) \times [-1, 1])$ contains at most two components. No two of these components are contained in the same component of $\partial D^3 \setminus \partial D$. Thus $\{\alpha(0), \alpha(1)\}$ is contained in a single component of $\partial D^3 \setminus p((\beta_1 \cup \beta_2) \times [-1, 1])$. Hence $\pi_1(p^{-1}(\alpha(0), \alpha(1)))$ is contained in a single component of $S^1 \setminus \{\beta_1 \cup \beta_2\}$. ■

Corollary 12 *If $|\hat{A}| \geq 4$ the bijection $h : \hat{A} \rightarrow \hat{A}$ is order preserving.*

Proof. Suppose $\gamma_1 < \gamma_2 < \gamma_3 < \gamma_4$ and $\gamma_i \in \hat{A}$. Let $int\phi X_i = \gamma_i$. Choose 4 points $x_i \in X_i$. Let $\lambda \subset X^*$ be an arc connecting x_1 to x_2 such that $int(\lambda) \subset U$. Let $\mathcal{A} \subset X^*$ be an arc connecting x_3 to x_4 such that $int(\mathcal{A}) \subset U$ and such that $\mathcal{A} \cap \lambda = \emptyset$. Let $\Lambda = \{\infty, -\infty\} \cup (\lambda \times (-1, 1)) \subset \overline{X^*} \times (-1, 1)$. Observe Λ is compact and homeomorphic to D^2 . Let $D = p\Phi(H^{**}\Lambda)$. Let $\alpha = p\Phi(H^{**}(\mathcal{A} \times \{0\}))$. Let $\beta_i = h(\gamma_i)$. Apply Lemma 11 to conclude β_3 and β_4 belong to the same component of $S^1 \setminus \{\beta_1 \cup \beta_2\}$. Thus by Lemma 7 h is order preserving. ■

8 Main results

Define $G(U) = \{H : \overline{U} \rightarrow \overline{U} \mid H \text{ is a homeomorphism and } H(U) = U\}$.

Define $G(U \times R) = \{H : \overline{U} \times R \rightarrow \overline{U} \times R \mid H \text{ is a homeomorphism and } H(U \times R) = U \times R\}$.

Recalling Theorems 1 and 5 define $rot(h) : G(U) \rightarrow R$ such that $rot(h)$ is the rotation number of $\overline{\psi h \psi^{-1}}_{S^1}$.

8.1 The case ∂U is not locally connected

Suppose ∂U is not locally connected.

Recall the bijection $h : \hat{A} \rightarrow \hat{A}$ from Corollary 10. Define a function $Rot : G(U \times R) \rightarrow R$ as follows.

If $|\hat{A}| \geq 3$ recall Corollary 12, note h is order preserving, and define $Rot(H) = Rot(g)$ where $g : S^1 \rightarrow S^1$ is any homeomorphism compatible with h .

If $|\hat{A}| \leq 2$ define $Rot(H) = 0$ if H is orientation reversing and fixes $\{\infty, -\infty\}$ pointwise or if H is orientation preserving and swaps

$\{\infty, -\infty\}$. Otherwise define $Rot(H)$ to be 0 or $\frac{1}{2}$ determined, respectively, by whether $h = ID_A$ or not.

8.2 The case ∂U is locally connected

The case ∂U is locally connected deserves special treatment since $\overline{U} \times R$ admits a homeomorphism extension theorem analogous to Theorem 1 and seems to provide a useful model of ‘3 dimensional prime end theory’ in the sense of [4].

Theorem 13 *Suppose ∂U is locally connected and $H \in G(U \times R)$. Then $\Psi H \Psi^{-1} : int(D^2) \times R \rightarrow int(D^2) \times R$ can be extended to a canonical homeomorphism of the 3 cell $(D^2 \times R) \cup \{\infty, -\infty\}$.*

Proof. Since ∂U is locally connected the conformal map $\psi^{-1} : int(D^2) \rightarrow U$ can be continuously extended to a surjective map $\psi^{-1} : D^2 \rightarrow \overline{U}$ (Theorem 2.1 p20 [10]). To show X^* is compact, it suffices, since X^* is complete, to show that each sequence in X^* has a Cauchy subsequence. Suppose y_n is a sequence in X^* . Let $x_n = \phi(y_n)$. Let $\{x_{n_k}\} \subset D^2$ be a Cauchy subsequence of $\{x_n\}$. Consider the chords $[x_{n_l}, x_{n_k}] \subset D^2$. Since ψ^{-1} is uniformly continuous, $\text{diam}(\psi^{-1}[x_{n_l}, x_{n_k}]) \rightarrow 0$. Hence $d^*(y_{n_l}, y_{n_k}) \rightarrow 0$. Consequently X^* is compact and the injective map from Lemma 2 $\phi : X^* \rightarrow D^2$ is a homeomorphism. Thus the homeomorphism from Theorem 9 $H^* : X^* \times R \rightarrow X^* \times R$ induces a homeomorphism of $D^2 \times R$ leaving invariant the ends $\{\infty\}$ and $\{-\infty\}$.

To see that H is canonical let $j : U^* \times R \rightarrow U \times R$ denote identity. Suppose $H_N \rightarrow H$ in $G(U \times R)$. Note $\{j^{-1}H_N j\}$ is uniformly equicontinuous and converges pointwise to $j^{-1}H j$. Hence $H_N^* \rightarrow H^*$ uniformly. Thus $\Psi H_N^* \Psi^{-1} \rightarrow \Psi H^* \Psi^{-1}$ uniformly. ■

To define rotation number when ∂U is locally connected we seek a nonempty proper totally disconnected set $B \subset S^1$ such that $B \times R$ is invariant under the induced homeomorphism $\overline{\Psi H \Psi^{-1}} : D^2 \times R \rightarrow D^2 \times R$.

The point $x \in \partial U$ is a **cutpoint** if $\partial U \setminus \{x\}$ not connected. Let $f : D^2 \rightarrow \overline{U}$ denote the continuous extension of ψ^{-1} . The map f is **light** since the prime ends of U are in bijective correspondence with points of S^1 (Theorem 2.15 [10]). Let $B = \{x \in S^1 \mid f(x) \text{ is a cutpoint of } \partial U\}$.

Lemma 14 $z \in B$ if and only if $|f^{-1}(z)| \geq 2$.

Proof. If $z \in \partial U$ and $|f^{-1}(z)| = 1$ then $z \notin B$ is since $S^1 \setminus f^{-1}(z)$ is connected and hence $f(S^1 \setminus f^{-1}(z)) = \partial U \setminus z$ is connected. ■

Suppose $|f^{-1}(z)| \geq 2$ choose $x \neq y$ such that $f(y) = f(x) = z$. Let $[x, y] \subset D^2$ denote the chord from x to y . If V is a component

of $\text{int}(D^2) \setminus [x, y]$ then $f(\partial V) \subset \partial(f(V))$. Thus, since f is light, each complementary domain of the simple closed curve $f([x, y])$ has nonempty intersection with ∂U . Hence $z \in B$.

Lemma 15 *Suppose $x \neq y$, $\{x, y\} \subset S^1$ and $z = f(x) = f(y)$. Suppose a and b lie in distinct components of $S^1 \setminus \{x, y\}$. Suppose $\{a, b\} \cap f^{-1}(z) = \emptyset$. Then $f(a) \neq f(b)$.*

Proof. The points a and b belong to the boundaries of distinct components of $\text{int}(D^2) \setminus [x, y]$. These components in turn map into distinct complementary domains of the simple closed curve $f([x, y])$, since points of $\text{int}(D^2)$ can approach the chord $[x, y]$ from distinct sides. By hypothesis $\{f(a), f(b)\} \cap f([x, y]) = \emptyset$. Thus $f(a)$ and $f(b)$ belong to distinct components of $f([x, y])$ and in particular $f(a) \neq f(b)$. ■

Lemma 16 *If ∂U is not a simple closed curve then $B \neq \emptyset$ and $B \neq S^1$.*

Proof. If ∂U is not a simple closed curve then by Lemma 14 f is not one to one and hence $B \neq \emptyset$. Let $\alpha_0 \subset S^1$ be any closed interval with distinct endpoints x_0 and y_0 such that $z_0 = f(x_0) = f(y_0)$. Proceeding recursively, if possible let $\alpha_{n+1} \subset \alpha_n \setminus f^{-1}(z_n)$ be a closed interval with distinct endpoints x_{n+1} and y_{n+1} such that $f(x_{n+1}) = f(y_{n+1})$ and $|x_{n+1} - y_{n+1}| < \frac{1}{n+1}$. Suppose the process never terminates. Let $a = \lim x_n = \lim y_n$. Suppose $f(b) = f(a)$. By Lemma 15 $b \in \cap \alpha_n$. Thus $b = a$ and by Lemma 14 $a \notin B$. If the process terminates then there must exist a nontrivial open interval $(x, y) \subset \alpha_0$ such that $f|_J$ is one to one and $f(x) = f(y)$. Let $a \in (x, y)$ and apply Lemma 15 to conclude $a \notin B$. ■

Lemma 17 *The following are equivalent. 1) $b \notin B$. 2) Suppose $g : S^1 \rightarrow \overline{U} \times R$ satisfies $g(1) = (f(b), \hat{t})$ and $g(\theta) \in U \times R$ if $\theta \neq 1$. Then there exists a map $G : D^2 \rightarrow \overline{U} \times R$ such that $G(z) \in U \times R$ whenever $z \in \text{int}(D^2)$ and $G_{S^1} = g$.*

Proof. Let $\pi_1 : \overline{U} \times R \rightarrow \overline{U}$ and $\pi_2 : \overline{U} \times R \rightarrow R$ denote the projection maps. Suppose $b \notin B$ and $g : S^1 \rightarrow \overline{U} \times R$ satisfies $\pi_1 g(1) = f(b)$ and $g(\theta) \in U \times R$ if $\theta \neq 1$. By Lemma 14 $|f^{-1}(f(b))| = 1$ and hence the formula $\alpha = f^{-1}(\pi_1 g) : S^1 \rightarrow D^2$ determines a map. Let $\beta : D^2 \rightarrow D^2$ be a continuous extension of α such that $\beta(z) \in \text{int}(D^2)$ if $z \in \text{int}(D^2)$. Define $G : D^2 \rightarrow \overline{U}$ via $G = f(\beta)$. Since R is contractible extend $\pi_2(g) : S^1 \rightarrow R$ to a map $h : D^2 \rightarrow R$. Define $G : D^2 \rightarrow \overline{U} \times R$ via $G(z) = (G(z), h(z))$.

Conversely suppose $b \in B$. By Lemma 14 choose $\hat{b} \in S^1$ such that $\hat{b} \neq b$ and $f(\hat{b}) = f(b)$. Let $[b, \hat{b}] \subset D^2$ denote the chord. Let $\hat{g}$

map S^1 homeomorphically onto the simple closed curve $f([b, b^\wedge]) \subset \overline{U}$ such that $g^\wedge(1) = f(b) = f(b^\wedge)$. Both components of $R^2 \setminus \text{im}(g^\wedge)$ contain points of ∂U . Hence $g^\wedge$ is essential in $\{f(b)\} \cup (R^2 \setminus \partial U)$. In particular G cannot exist since $\pi_1(G)$ would show that $g^\wedge$ is inessential. ■

Since condition 2) is a topological property $B \times R$ maps onto itself under the induced homeomorphism $\overline{\Psi H \Psi^{-1}} : D^2 \times R \rightarrow D^2 \times R$. It follows that if $B^\wedge = \{x \in S^1 \mid \{x\} \text{ is a component of } B \text{ or } x \text{ is an endpoint of a nontrivial interval of } B\}$ then $B^\wedge \neq \emptyset$, $B^\wedge$ is totally disconnected, and $\overline{\Psi H \Psi^{-1}}(B^\wedge \times R) = B^\wedge \times R$. If $|B^\wedge| \neq 2$ let $g : S^1 \rightarrow S^1$ be any homeomorphism such that $g(x) = y$ whenever x and y are components of $B^\wedge$ such that $\overline{\Psi H \Psi^{-1}}(\{x\} \times R) = \{y\} \times R$. Define $\text{Rot}(H) = \text{rot}(g)$. If $|B^\wedge| = 2$ define $\text{Rot}(H) = 0$ if $\overline{\Psi H \Psi^{-1}}$ is orientation reversing and fixes $\{\infty, -\infty\}$ pointwise or if $\overline{\Psi H \Psi^{-1}}$ is orientation preserving and swaps $\{\infty, -\infty\}$. Otherwise define $\text{Rot}(H)$ to be 0 or $\frac{1}{2}$ determined, respectively, by whether $\overline{\Psi H \Psi^{-1}}$ interchanges the lines of $B^\wedge \times R$ or not.

8.3 Main Theorem on rotation number over $U \times R$

Theorem 18 *Suppose ∂U is not a simple closed curve. The function $\text{Rot} : G(\overline{U} \times R) \rightarrow R$ is continuous. If $F, H \in G(\overline{U} \times R)$ then $\text{Rot}(H) = \text{Rot}(FHF^{-1})$. If there exists $h \in G(\overline{U})$ such that $\forall z, t \ H(z, t) = (h(z), t)$ then $\text{Rot}(H) = \text{rot}(h)$.*

Proof. If $|A^\wedge| \geq 3$ then by Corollary 12 Rot is well defined. Suppose $g, f : S^1 \rightarrow S^1$ are homeomorphisms compatible, respectively with H and F . Then fgf^{-1} is compatible with FGF^{-1} . Hence $\text{Rot}(H) = \text{rot}(g) = \text{rot}(fgf^{-1}) = \text{Rot}(FGF^{-1})$. For continuity of Rot suppose $H_n \rightarrow H$. Let $h_n : A^\wedge \rightarrow A^\wedge$ and $h : A^\wedge \rightarrow A^\wedge$ denote the corresponding order preserving bijections. If J is a nontrivial component of $A^\wedge$ then $h_n(J)$ is eventually constant. Construct compatible homeomorphisms $g_n : S^1 \rightarrow S^1$ and $g : S^1 \rightarrow S^1$ as in Lemma 7. Note eventually $g_n|_J = g|_J$ and hence $\text{rot}(g_n) = \text{rot}(g)$ eventually. If each component J of $A^\wedge$ is trivial then $g_n \rightarrow g$ pointwise on a dense set $A^\wedge \subset S^1$ and hence $g_n \rightarrow g$ uniformly. Thus by Theorem 5 $\text{rot}(g_n) \rightarrow \text{rot}(g)$.

If ∂U is not locally connected and $|A^\wedge| \leq 2$ the theorem follows from the definition of Rot .

If ∂U is locally connected the homeomorphism $\overline{\Psi H \Psi^{-1}}$ varies continuously with H . The theorem follows from the definition of Rot and from Theorem 5. ■

References

- [1] Alligood, Kathleen T.; Yorke, James A. Fractal basin boundaries and chaotic attractors. *Chaos and fractals* (Providence, RI, 1988), 41–55, *Proc. Sympos. Appl. Math.*, 39, Amer. Math. Soc., Providence, RI, 1989.
- [2] Barge, Marcy; Gillette, Richard M. Indecomposability and dynamics of invariant plane separating continua. *Continuum theory and dynamical systems* (Arcata, CA, 1989), 13–38, *Contemp. Math.*, 117, Amer. Math. Soc., Providence, RI, 1991.
- [3] Barge, Marcy Prime end rotation numbers associated with the Hénon maps. *Continuum theory and dynamical systems*, 15–33, *Lecture Notes in Pure and Appl. Math.*, 149, Dekker, New York, 1993.
- [4] Brechner, Beverly L.; Lee, Joo S. A three-dimensional prime end theory. *Topology Proc.* 20 (1995), 15–47.
- [5] Cartwright, M. L.; Littlewood, J. E. Some fixed point theorems. *Ann. of Math.* (2) 54, (1951). 1–37.
- [6] Devaney, Robert L. *An introduction to chaotic dynamical systems*. The Benjamin/Cummings Publishing Co., Inc., Menlo Park, CA, 1986.
- [7] Fabel, Paul Self-homeomorphisms of the 2-sphere which fix pointwise a nonseparating continuum. *Fund. Math.* 155 (1998), no. 3, 201–214.
- [8] Franks, John Rotation numbers and instabilty sets. *Bull. Amer. Math. Soc. (N.S.)* 40 (2003), no. 3, 263–279 (electronic)
- [9] Mayer, John C.; Oversteegen, Lex G. Denjoy meets rotation on an indecomposable cofrontier. *Continuum theory and dynamical systems*, 183–199, *Lecture Notes in Pure and Appl. Math.*, 149, Dekker, New York, 1993.
- [10] Pommerenke, Ch. Boundary behaviour of conformal maps. *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, 299. Springer-Verlag, Berlin, 1992.
- [11] Ursell, H. D.; Young, L. C. Remarks on the theory of prime ends. *Mem. Amer. Math. Soc.*(1951). no. 3, 29 pp.